



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DIOPHANTINE ANALYSIS.

70. Proposed by CHARLES CARROLL CROSS, Libertytown, Md.

Give methods for decomposing numbers into squares, cubes, or biquadates and show that 61×200^3 is the sum of ten cube numbers and that 844933 is the sum of eleven biquadrates in thirteen different ways. [From *The Mathematical Magazine*, Vol. II, No. 10.]

Solution by the PROPOSER.

Let s_1 = sum of products taking 1 number at a time(1),
 s_2 = sum of products taking 2 numbers at a time.....(2),
 s_3 = sum of products taking 3 numbers at a time.....(3),
.....
 s_n = sum of products taking n numbers at a time.....(n).
Then $a + b + c + d \dots n = s_1 \dots (b_1)$,
 $(1)(b_1) - 2(2)$ gives
 $a^2 + b^2 + c^2 + d^2 \dots n^2 = s_1^2 - 2s_2 \dots (b_2)$,
 $(1)(b_2) - [(2)(b_1) - 3(3)]$ gives
 $a^3 + b^3 + c^3 + d^3 \dots n^3 = s_1^3 - 3s_1s_2 + 3s_3 \dots (b_3)$,
 $(1)(b_3) - [(2)(b_2) - (3)(b_1) - 4(4)]$ gives
 $a^4 + b^4 + c^4 + d^4 \dots n^4 = s_1^4 - 4s_1^2s_2 + 4s_1s_3 + 2s_2^2 + 4s_4 \dots (b_4)$,
 $(1)(b_4) - [(2)(b_3) - (3)(b_2) - (4)(b_1) - 5(5)]$ gives
 $a^5 + b^5 + c^5 + d^5 \dots n^5 = s_1^5 - 5s_1^3s_2 + 5s_1^2s_3 + 5s_1s_2^2 + 5s_1s_4 - 5s_2s_3 + 5s_5 \dots (b_5)$,
etc. etc. etc.

[In (b_5) let $s_1 = x$, $s_2 = -xy$, $s_3 = 2xy^2$, $s_4 = y^2(x^2 - y^2)$, and $s_5 = \frac{1}{5}y^5$, then $a^5 + b^5 + c^5 + d^5 + e^5 + f^5 = (x + y)^5$. This answers Dr. Drummond's note on page 182, Vol. V., of the MONTHLY.]

71. Proposed by A. H. BELL, Hillsboro, Ill.

Find five numbers such that the product of any two plus 1 will equal a square.

I. Partial Solution by CHARLES C. CROSS, Libertytown, Md.

We readily find four numbers to be $(n-1)$, $(n+1)$, $4n$, and $4n(4n^2-1)$.

Let $n=2, 3, 4$, and so on, and we get

1, 3, 8, 120,	also	1, 8, 15, 528,
2, 4, 12, 420,		2, 12, 24, 2380,
3, 5, 16, 1008,		3, 16, 33, 6440,
4, 6, 20, 1980,		4, 20, 42, 13572,
etc., etc.		etc., etc.

From which we get a more general formula for four numbers and find it to be, m , $n^2-1+(m-1)(n-1)^2$, $n(mn+2)$, $m(m^2n-2mn+mn^2+4n-2)^2+(m^2n-2mn+mn^2+4n-2)$.

If there can be found a value for n that will render $(n-1)(128n^3-8n-8)$
 $(256n^6-256n^4+63n^2-2n-1)+(256n^6-288n^4-32n^3+65n^2+2n+1)^2=\square$ then
 the five numbers are $(n-1)$, $(n+1)$, $4n$, $4n(4n^2-1)$, and $[(256n^6-256n^4+63n^2-2n-1)(128n^3-8n-8)]/(256n^6-288n^4-32n^3+65n^2+2n+1)^2$; but life is too
 short to attempt this.

Dr. Zerr says, "I have not been able to find the five numbers."

If you get no answers to problem 71, I can give Legendre's (improved),
 and it is a fractional answer for the fifth number. Mr. Wilkes has found four
 numbers, as follows: $n-1$, $n+1$, $4n$, $16n^3-4n$. Taking these for four of them
 then I can prove that if there is another integral number it must end in 0.

I have investigated, up to numbers having over five thousand digits, with-
 out finding any to answer.

A. H. BELL.

I have given considerable time to problem 71, but as yet have failed to
 obtain five numbers. I have found general values for three such numbers, viz.,
 m , $n(mn\pm 2)$, and $(n+1)[(n+1)m\pm 2]$. I have also found partial general values
 for four such numbers, viz., m , $m\pm 2$, $4(m\pm 1)$, and $16(m\pm 1)^3-4(m\pm 1)$; but one
 of the sets of four numbers that I obtained by inspection, can not be obtained by
 this formula. The set is 1, 8, 15, 528.

M. A. GRUBER.

MISCELLANEOUS.

63. Proposed by F. P. MATZ, D. Sc., Ph. D., Professor of Mathematics and Astronomy, Irving College, Me-
 chanicsburg, Pa.

Show that the path of a projectile moving with a constant velocity is an inverted
 catenary of equal strength.

Solution by GEORGE R. DEAN, Professor of Mathematics, University of Missouri School of Mines and Metal-
 lurgy, Rolla, Mo.

The differential equations are, obviously,

$$\frac{d^2y}{dt^2} = -g, \quad \frac{ds}{dt} = c.$$

$$\text{Whence, } \frac{dy}{dt} = -gt + k, \quad \frac{dx}{dt} = \sqrt{c^2 - (gt - k)^2}, \quad x + k' = -\frac{1}{g} \cos^{-1} \frac{gt - k}{c}.$$

$$\text{By division, } \frac{dy}{dx} = \frac{-\cos g(x + k')}{\sqrt{c^2 - c^2 \cos^2 g(x + k')}} = -\cot g(x + k'),$$

$$\text{or, } \cot^{-1} \frac{dy}{dx} = -g(x + k').$$

Differentiation gives